

The Distances to Open Clusters as Derived from Main-sequence Fitting. II. Construction of Empirically Calibrated Isochrones¹

Marc H. Pinsonneault and Donald M. Terndrup

Department of Astronomy, The Ohio State University, Columbus, OH 43210

`pinsono,terndrup@astronomy.ohio-state.edu`

Robert B. Hanson

University of California Observatories/Lick Observatory, Santa Cruz, CA 95064

`hanson@ucolick.org`

and

John R. Stauffer

*Infrared Processing and Analysis Center, California Institute of Technology, Mail Code 100-22,
770 South Wilson Avenue, Pasadena, CA 91125*

`stauffer@ipac.caltech.edu`

ABSTRACT

We continue our series of papers on open cluster distances by comparing multicolor photometry of single stars in the Hyades to theoretical isochrones constructed with various color-temperature relations. After verifying that the isochrone effective temperatures agree well with spectroscopically determined values, we argue that mismatches between the photometry and the theoretical colors likely arise from systematic errors in the color-temperature relations. We then describe a method for empirically correcting the isochrones to match the photometry, and discuss the dependence of the isochrone luminosity on metallicity.

Subject headings: stars: distances, stars: abundances, stars: evolution, open clusters: individual (Hyades)

¹This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

1. Introduction

Open clusters are important laboratories for testing stellar evolution models and for deciphering the star formation history of the Galaxy, since each cluster contains samples of stars of a single age and (probably) composition. As is well known, determining distances is the most fundamental step to measuring cluster ages and other key properties.

The success of the Hipparcos mission (ESA 1997) has now provided us with many cluster distances derived from stellar parallaxes. While the new astrometry is a significant advance, it did not come without some controversy. In particular, there were a few cases, notably the Pleiades, in which the Hipparcos parallax distance (Mermilliod et al. 1997; Robichon et al. 1999; van Leeuwen 1999) was significantly different from that obtained by main-sequence (MS) fitting using stellar isochrones (Pinsonneault et al. 1998; Soderblom et al. 1998). A number of papers have discussed these discrepancies in cluster distance estimates, which could have several causes. First, the systematic or random errors in MS-fitting distances could be significantly larger than estimated, as claimed by Robichon et al. (2000), or the photometry, reddening, and/or metal abundances in the best-studied clusters could be seriously in error. In the case of the Pleiades, the cluster may have an unusually high helium abundance, making its main sequence ≈ 0.3 mag fainter than would be expected from its metallicity (Belikov et al. 1998); this, however, is difficult to understand since there do not seem to be nearby field stars of similar characteristics in the Hipparcos catalog (Soderblom et al. 1998). Alternatively, the metal abundance from spectroscopy may have been overestimated (Percival, Salaris, & Kilkenney 2003, and references therein); this point has been disputed by, *inter alia*, Stello & Nissen (2001). Second, there could be some overlooked flaw in the stellar models used to compute isochrones or in the transformation of model quantities to broadband colors. Particularly on the lower main-sequence in young clusters, stellar activity producing spots or chromospheric emission can affect colors in the blue (van Leeuwen, Alphenaar, & Meys 1987; Stauffer et al. 2003). Finally, the cluster parallax discrepancies, amounting to ~ 1 mas, may just reflect the actual size of Hipparcos’ parallax errors over the small angular scale ($\lesssim 1^\circ$) subtended by most open clusters except the Hyades. On scales comparable to Hipparcos’ $0^\circ.9$ field of view, the individual stellar parallaxes are correlated (van Leeuwen & Evans 1998), causing local zero-point errors in the cluster parallaxes. These have been suggested to be ~ 1 mas in the Pleiades (Makarov 2002) and Coma Ber (Makarov 2003, preprint), but smaller in the Hyades (Narayanan & Gould 1999a,b; de Bruijne, Hoogerwerf, & de Zeeuw 2001). Recently, there has been a similar discussion about the distance to NGC 2516 (Terndrup et al. 2002; Percival, Salaris, & Kilkenney 2003); here, uncertainties in the cluster metallicity and in the amount of foreground extinction complicate the comparison between the MS-fitting distance and the Hipparcos measurement. Regardless of the source or sources of these discrepancies, it remains the case that the 1997-1999 analysis of Hipparcos parallaxes and current stellar evolutionary models are not mutually consistent when applied to the ensemble of nearby clusters with precise parallaxes.

The idea that MS-fitting can be used to find distances to clusters and even individual stars has, of course, been around for a long time (e.g., Eggen 1948; Johnson & Knuckles 1955), but it

has proven difficult to develop a precise technique using theoretical isochrones, even for determining relative distances². The stellar evolution models themselves are now well constrained in the case of the Sun (e.g., Basu, Pinsonneault, & Bahcall 2000; Bahcall, Pinsonneault, & Basu 2001), but are less certain for the full range of temperatures and luminosities in any open cluster. In order to generate isochrones, the models need transformations from theoretical quantities (L, T_{eff}) into observational magnitudes and colors. A number of different color-temperature calibrations are available, including some generated empirically (e.g., Gratton, Carretta, & Castelli 1996; Weiss & Salaris 1999; Lejeune & Schaerer 2001; VandenBerg & Clem 2003). Despite improvements in the determination of atmospheric opacities, bolometric corrections, etc., there typically remain small but significant mismatches between calculated and isochrone colors and the best photometry in open clusters (e.g., Terndrup et al. 2000; de Bruijne, Hoogerwerf, & de Zeeuw 2001; Castellani, Degl’Innocenti, & Prada Moroni 2001; Grocholski & Sarajedini 2003). The size of these discrepancies can be as large as 0.04 – 0.06 mag in broadband colors. Furthermore, the dependence on metallicity of the absolute magnitude M_V of the MS is a function of color, being greater when bluer colors such as $B - V$ are used in the color-magnitude diagram (CMD) and less, for example, when $V - I$ or $V - K$ are employed. Taken together, all this means that the derived distance depends on the selection of colors used in the MS-fitting (Pinsonneault et al. 1998). On the other hand, if the isochrones can be accurately calibrated over a wide range of colors, it may be possible to use multiband photometry to derive more accurate distance and reddening estimates and to provide a photometric abundance indicator for open clusters.

Motivated by the continuing discussion of open cluster distances, we have undertaken a careful reexamination of all the ingredients of the MS-fitting technique, revising and extending our previous formulation (Pinsonneault et al. 1998) and taking advantage of new calibrations from Hipparcos and from improvements to our stellar evolution models. Our first paper (Pinsonneault et al. 2003, hereafter Paper I) discussed the construction of the isochrones and demonstrated that they provided a good match to the mass/luminosity relation for the components of the eclipsing binary vB 22 in the Hyades. In this paper, we perform additional external checks on the metallicity and temperature scales adopted for the Hyades, then use photometry to generate empirical adjustments to the isochrone colors and bolometric corrections. We conclude by discussing the sensitivity of the MS luminosity to metallicity for various available color calibrations. In subsequent papers of this series, we will extend the calibration to fainter magnitudes, provide a grid of calibrated isochrones appropriate for open clusters, and determine the effects of random and systematic errors in the main-sequence fitting method.

²Alternative methods using field stars with measured parallaxes have equivalent challenges; see, for example, VandenBerg et al. (2000).

2. The empirical calibration

2.1. Data

We adopted the Hyades as the zero point for the MS-fitting method for a number of reasons. The cluster has a negligible reddening $E(B - V) = 0.003 \pm 0.002$ (Crawford 1975; Taylor 1980) and excellent membership information from proper motions and radial velocities. Although the cluster’s size is a significant fraction of its distance, the Hipparcos parallaxes and proper motions allow the determination of the distances to individual Hyades members to a precision of $\sim 2\%$ (de Bruijne, Hoogerwerf, & de Zeeuw 2001), with reasonable assumptions about the cluster’s internal motions. The average distance, taken as that of the cluster’s center of mass ($m - M = 3.33 \pm 0.01$, or $d = 46.34 \pm 0.27$ pc) from Perryman et al. (1998), is not altered significantly by correlations of the Hipparcos parallaxes on small angular scales (Narayanan & Gould 1999b; de Bruijne, Hoogerwerf, & de Zeeuw 2001). The most significant problem is that the Hipparcos data do not extend below about $M_V = 8$, insufficient for calibrating isochrones for the lower MS. This will be corrected by making use of data in other clusters such as the Pleiades or Praesepe, as discussed in a future paper.

We chose as our Hyades sample a set of stars identified as members of the cluster by Perryman et al. (1998), excluding any star that is a known or suspected binary. Most of these are included in the high-resolution spectroscopic sample of Paulson, Sneden, & Cochran (2003, hereafter PSC).

To construct a calibrating set of photometry for the Hyades, we inspected the numerous UBV photometric studies and selected those which were in mutual agreement. For B and V , we selected stars from Johnson & Knuckles (1955), Mendoza (1967), and the Hipparcos mission (ESA 1997). In V , we also had photometry from Weis and collaborators (Ugoren & Weis 1977; Weis, Delucca, & Ugoren 1979) that was also used for (Kron) $V - I_K$ colors (these will be discussed below). The Hipparcos catalog provides individual errors for stars, and the quoted errors in the Johnson and Mendoza papers respectively are 0.004 and 0.01 magnitudes. The dispersion in $B - V$ between these three samples is consistent with the quoted errors, and the zero-point difference for stars in common is negligible (of order 0.001 mag or less).

The Hipparcos catalog lists V magnitudes for all of the stars in our sample; all but 32 of these are derived from ground-based photometry. There is a small (0.012 mag) zero-point offset between the space-based Hipparcos V magnitudes and the Johnson/Mendoza data, with dispersions of 0.013 and 0.009 mag respectively; the difference is in the sense that the ground-based data are fainter. There are only six stars in common between Weis and Johnson, with a formal dispersion of 0.015 mag and a zero-point offset of 0.005 (Johnson fainter), which is not statistically significant. We choose the ground-based data for our zero-point, adding 0.012 mag to the Hipparcos space-based V magnitudes, and averaged the results assuming an error of 0.01 for Hipparcos, Mendoza, and Johnson and 0.015 for Weis.

The apparent V magnitudes were converted into absolute M_V magnitudes using the individual

distance moduli from de Bruijne, Hoogerwerf, & de Zeeuw (2001). The errors for M_V are larger than for V (or V used in the construction of $V - K_s$ colors), since the error in the distances are included and these errors are usually larger than those in V alone.

We used two sources for the K band data: Carney (1982), which is on the CTIO-CIT system (Frogel et al. 1978; Elias et al. 1982), and data from the 2MASS All Sky release. The latter is on the “short- K ” (K_s) system (Persson et al. 1998; Carpenter 2001). We converted the Carney (1982) K magnitudes to K_s using the transformation in Carpenter (2001) and averaged these with the 2MASS values, using Carney’s error estimate of 0.011 mag for each star. The error in the $V - K$ color was obtained by adding the errors in V and K_s in quadrature.

The merger of the I -band data for the clusters was somewhat more involved than for BVK , largely because of the need to transform different I -band systems (Johnson, Kron/Eggen, and Cousins) onto the Cousins system. There were three main sources of Johnson data with a total of 60 stars: Mendoza (1967), Johnson et al. (1966), and Johnson, MacArthur, & Mitchell (1968). In addition there were six stars from Carney & Aaronson (1979).

Weis and collaborators (Upgren & Weis 1977; Weis, Delucca, & Upgren 1979; Weis & Upgren 1982; Upgren, Weis, & Hanson 1985; Weis & Hanson 1988) have $V - I$ data in the Kron system for 23 stars. There were also 11 stars with Johnson $V - I$ and accurate $R - I$ colors from Eggen (1982), which are on the Kron system (Eggen 1975); we found that although the Eggen and Johnson $R - I$ colors could be accurately transformed onto the same system, the $V - I$ colors from Eggen had larger errors. This is probably because Eggen’s V magnitudes were not obtained at the same time as $R - I$, so the errors in $V - I$ are much greater than in $R - I$. Only five stars in the sample have $V - I$ colors in both the Kron and Johnson systems. We were unable to achieve a good transformation between the merged colors and those in Taylor & Joner (1985), so we ignored their measurements.

The Johnson data were transformed onto the Cousins system using the prescription in Bessel (1979), while the Weis and Eggen data were transformed onto the Cousins system using the cubic formula in Bessel & Weis (1987). The average difference between the transformed Johnson and Weis/Eggen colors are 0.009 and 0.004 mag respectively, with a dispersion of 0.012 and 0.013 mag. These error estimates are only approximate because the number of stars with multiple $V - I$ errors was small: the bright ($V \lesssim 8$) stars in the Hyades mostly have Johnson photometry while the faint stars have Kron photometry and the magnitude range of overlap is limited. We therefore assumed that the colors on both systems were correctly transformed onto the Cousins system, and did not apply any further zero-point or slope offsets to the transformed $V - I$ data.

The list of merged photometry is presented as Table 1. The Hipparcos designation for each star is in the first column, while the second lists other names. The remaining columns show the visual magnitude and colors, where the errors represent either quoted errors in the various datasets if only one measurement is available, or the standard deviation if more than one datum was averaged.

For completeness, we have also included the $J - K$ and $H - K$ colors for each star, even though

they are not otherwise discussed in this paper (in particular, the metallicity sensitivity of $J - K$ is smaller than for $V - K$, and there is very little variation in the values of $H - K$ for main-sequence stars). Note that several of the brightest stars have large errors in the infrared colors because they are saturated in the 2MASS survey. We plan to discuss the full set of infrared colors and their usefulness for reddening and metallicity determinations in a subsequent paper.

2.2. Comparing different color calibrations

In Paper I, we described the construction of an isochrone for the Hyades and demonstrated that it compared favorably to the masses and luminosities of the components of the Hyades eclipsing binary vB 22. The parameters for the Hyades depend somewhat on the helium abundance assumed for the Hyades and on the adopted solar model used to calibrate the mixing length parameter for convection. The isochrone used here is the same as in Paper I, namely constructed from models ignoring microscopic diffusion with $Y = 0.273$ and $\alpha = 1.72$. The age of the Hyades was taken as 550 Myr, appropriate for models lacking convective overshoot (e.g., Perryman et al. 1998). The models used the solar abundance mix of Grevesse & Noels (1993), and scaled to a Hyades abundance of $[\text{Fe}/\text{H}] = +0.13 \pm 0.01$ (PSC), using the Sun’s relative abundances. The PSC average abundance is quite close to that found by Boesgaard & Friel (1990); see Perryman et al. (1998) for a summary of previous abundance estimates in the Hyades. The input physics is similar to that used in the recent Yi, Kim, & Demarque (2003) Y^2 isochrones for stars of solar mass and above. Our usage of the Saumon, Chabrier, & Van Horn (1995) equation of state for lower mass main sequence stars does cause intrinsic differences between our isochrones and theirs for stars below 0.8 solar masses.

In Figures 1–3 we compare the theoretical isochrone to the Hyades photometry in Table 1, showing the effect of using several available color-temperature calibrations. Our base case, displayed as the solid line in these figures, uses the corrected color-temperature relation in Lejeune et al. (1998), which employed multicolor data to adjust theoretical flux distributions based on Kurucz and Allard & Hauschildt (1995) atmospheres. This isochrone matches the photometry reasonably well, although systematic departures from the data are readily apparent especially for the reddest stars.³ The three color-magnitude diagrams (CMDs) are in M_V and $B - V$, $V - I_C$, and $V - K_s$, respectively. The absolute magnitudes were generated from the V magnitudes and individual kinematic parallaxes from de Bruijne, Hoogerwerf, & de Zeeuw (2001) adjusted to a common distance modulus of $m - M = 3.33 \pm 0.01$ (Perryman et al. 1998).

Also shown on Figures 1–3 as a short-dashed line is the same isochrone generated with the color-temperature relation in Alonso et al. (1995, 1996); equivalently, the long-dashed line displays the isochrone made with the Lejeune et al. (1998) calibration *before* their application of empirical

³The various available color-temperature relations differ the most for stars cooler than about 4500 K. The empirically corrected Lejeune et al. (1998) relation, for example, is nearly identical to that presented by Flower (1996), but is bluer by nearly 0.1 mag at 4000 K.

corrections to the theoretical colors. Here, we will refer to the latter as the uncorrected Lejeune et al. (1998) calibration, though note that their terminology is different.

2.3. Isochrone effective temperatures

We have just shown that a single theoretical isochrone will produce different loci in the color-magnitude diagram when different color-temperature relations are applied. Before correcting the colors to match the Hyades photometry, it is necessary to do one additional test: namely to verify that the theoretical quantities generated by the stellar evolution models (L , T_{eff} , $\log g$) are reasonably close to that of the actual cluster, since any errors will be washed away by forcing the isochrone colors to match the photometry. The recent spectroscopic temperature determinations by PSC allow this test to be done with precision.

In both theoretical models and direct parallax measurements, high metallicity stars appear fainter at a fixed color than low metallicity stars. This is partially a stellar interiors effect; increased metal abundance makes stars of fixed mass slightly fainter and significantly cooler, moving them above the MS locus of lower-metallicity stars. Increased line blanketing in more metal-rich stars also makes them appear redder at fixed effective temperature. Increasing the helium abundance will make stars appear systematically fainter at fixed effective temperature or color; however, the spectral energy distribution and color-temperature relationship is largely insensitive to the helium abundance.

We now proceed to examine the match between the temperature and gravities in the isochrone to parameters derived by PSC. They determined effective temperatures for each star by the requirement that the Fe I abundances from several lines were independent of excitation potential. Gravity was found from requiring ionization equilibrium that yields the same abundance for Fe I and Fe II lines.

In Figure 4, we display (top panel) the difference between the PSC temperatures and those from the isochrone with the Lejeune et al. (1998) color calibration, namely the isochrone plotted as a solid line in Fig. 3. The difference is in the sense of isochrone *minus* spectra, and the isochrone values for each star were found by looking up the effective temperature at each M_V . These temperatures are therefore independent of systematic errors in the color calibration, discussed below in § 3. The lower panel shows the difference in $\log g$ constructed in a similar manner. The circular points show stars that are in our Table 1, while the triangles show other stars in PSC; these were not included in our sample either because they may be binaries or because there was accurate photometry but not accurate luminosities.

Overall, the agreement between the isochrone parameters and the spectroscopically derived values is remarkably good, with average differences of only -24 K in T_{eff} and $+0.02$ in $\log g$. There are, however, noticeable trends with temperature, in that the PSC temperatures are higher than the isochrone values for the hottest stars in their sample; for these stars, their adopted gravities are

somewhat too low. In combination, their temperatures and gravities imply masses about $0.2M_{\odot}$ above the isochrone values, or radii that are about 5% too large at fixed mass compared to the isochrones.

In their Table 7, PSC provide coefficients showing the sensitivity of the derived abundances to changes in temperature and gravity, from which we can rescale their $[\text{Fe}/\text{H}]$ to the temperatures and gravities of the isochrones (we adopt their microturbulence values without change). The coefficients were tabulated for two stars at the extremes of the temperature distribution, from which we linearly interpolated to determine the sensitivity of the abundances at any intermediate temperature. Figure 5 displays (top panel) the PSC values of $[\text{Fe}/\text{H}]$ as a function of isochrone effective temperature, where the symbols are the same as in the previous figure. The rescaled abundances are displayed in the lower panel.

The mean derived spectroscopic abundance using the rescaled abundances is $\langle[\text{Fe}/\text{H}]\rangle = +0.12 \pm 0.01$. This is in excellent agreement with the original assumed abundance of $+0.13$, showing the locus of the isochrone in the theoretical plane is fully consistent with spectroscopic abundance data to a high degree of accuracy.⁴ The dispersion in the rescaled abundances is slightly smaller than that obtained with the spectroscopic temperatures and gravities, the number of outliers is reduced, and any underlying trends in $[\text{Fe}/\text{H}]$ with T_{eff} are small. We therefore conclude that the models in the theoretical plane are a reasonable representation of the actual cluster.

2.4. A recap

To summarize, our argument so far has been thus: (1) Solar models are in excellent agreement with the stringent tests possible with helioseismic data (Basu, Pinsonneault, & Bahcall 2000), so models for Hyades stars of similar effective temperatures should be similarly well constrained; (2) The theoretical mass/luminosity relationship is in good agreement with the values obtained from the eclipsing binary vB 22 for a reasonable Hyades helium abundance (paper I); (3) Available color-temperature relations applied to the same isochrone do not match the photometry of the Hyades (§ 2.2); (4) The spectroscopic temperatures of Hyades are consistent with the luminosity / effective temperature relationship in the models (§ 2.3).

We therefore believe that the isochrones in the theoretical luminosity-effective temperature plane are well matched to the Hyades. In other words, the stellar interiors models are not the cause of the mismatch in shape between the isochrones and the Hyades photometry in Figs. 1–3. The deviations for any one isochrone are usually about the same size as the differences between

⁴Even though we started with the PSC value in constructing the isochrone, the agreement we find does not amount to a circular argument. If we had adopted (say) $[\text{Fe}/\text{H}] = 0.0$ for the isochrone, that isochrone would have been too faint at the Hipparcos distance to the cluster. The temperatures derived from M_V would be therefore be too high, which would have required the PSC abundances to be increased significantly.

published color calibrations, which suggests that we would be justified in applying modest empirical corrections to the isochrones.

We now proceed to define the empirical locus of the Hyades in the M_V -color plane and obtain a set of corrections to our base color-temperature relationship (Lejeune et al. 1998). We begin by examining the internal consistency of effective temperatures derived from published color-temperature relationships for the same star but with multiple colors. We then define empirical color-color relationships for the Hyades, tying the other colors to $B - V$, and identifying the locus of the cluster in the $(M_V, B - V)$ plane. When combined with the color-color relationships, this defines the locus of the cluster in each of the color indices that we have included in this paper (namely $B - V$, $V - I_C$, and $V - K_s$; other colors will be treated in subsequent papers). Finally, we use the isochrone M_V, T_{eff} relationship to convert these empirical corrections as a function of M_V into empirical corrections to the color-temperature relationship.

2.5. The internal consistency of the color-temperature relationships

If the metallicity and reddening of a cluster are known, a self-consistent set of color-temperature relationships should yield the same temperatures for each star regardless of which color is used to derive the temperature. In Figure 6, we compare the T_{eff} estimates for each star in Table 1 using the Lejeune et al. (1998) color calibration. The horizontal axis displays the isochrone effective temperature derived from M_V ; note that the temperatures extend to higher values than in Figures 4 and 5 because the PSC survey only included stars cooler than about $T_{\text{eff}} = 6300$ K. The top panel shows the difference between the temperature estimated from $V - I_C$ and that from $B - V$; the middle and lower panels, respectively, compare the temperature from $V - K_s$ to $B - V$ and $V - I_C$, again as a function of the temperature from M_V . Errors in the temperatures are propagated from errors in the photometry.

While the mean temperature differences are nearly zero (the largest is $\langle \Delta T \rangle = -60$ K for the data in the top panel), there are clear systematic differences as a function of temperature which can be as large as 300 K. This indicates that there are temperature ranges where the Lejeune et al. (1998) color calibration is not internally consistent, at least with respect to the Hyades multicolor photometry. These small inconsistencies may have arisen because the color transformation was derived by comparison to an ensemble of field stars with various metallicities; the (generally) small differences noted here would probably not have been apparent in the Lejeune et al. analysis.

We also explored internal inconsistencies using other color-temperature relationships, and found similar results. We therefore conclude that the color-temperature calibrations need to be modified not only to remove systematic errors in distances and photometric metallicity, but also for internal consistency. Recently, Grocholski & Sarajedini (2003) have compared multicolor photometry in many clusters to a number of color-temperature relations and have come to a similar conclusion.

2.6. Correcting the color-color relationship

The next step is to correct the color-color relationship. There were several reasons why we performed this correction rather than adjusting the isochrones in the (M_V, color) planes individually. First, the errors in the colors are typically smaller than the errors in M_V ; the latter, of course, arise from parallax errors and are much larger than the error in the V magnitudes. Second, correcting the color-color relation necessarily produces the desired result that the isochrone color-temperature relation be internally consistent (i.e., each color yields the same temperature for a star, at least to the limits set by photometric errors).

The smallest observational errors are in $B - V$, so we chose to define corrections in the other colors as a function of $B - V$. The method is illustrated in Figure 7. The top panel shows the color difference $\Delta(V - I_C)$ between the isochrone and the photometry for each star in Table 1, plotted against the observed $B - V$ color. The lower panel shows the same, but for $V - K_s$. The isochrone employed was the same base case (above), namely for $[\text{Fe}/\text{H}] = +0.13$ and the Lejeune et al. (1998) color calibration; we have neglected systematic errors arising from the small uncertainty in the metallicity of the Hyades. The color differences are in the sense of (photometry – isochrone), so positive values indicate that the star is redder than the isochrone. The isochrone colors $V - I_C$ and $V - K_s$ were computed by finding the effective temperature at each $B - V$, then looking up the other colors at that temperature. The error bars on each point include both the observational error in the photometry and the error in the predicted color arising from errors in $B - V$.

The smooth line in each panel of Figure 7 represents the color correction to $V - I_C$ or $V - K_s$ as a function of $B - V$ (specifically the $B - V$ color of the Hyades stars). The line was constructed by finding the average value of $\Delta(V - I_C)$ or $\Delta(V - K_s)$ in a moving box containing 3–5 points sorted by increasing $B - V$. The line was then smoothed by averaging each point with the linear interpolation of adjacent points blueward and redward.

These corrections would generate $V - I_C$ and $V - K_s$ colors that are consistent with the Hyades photometry once the isochrone generates the observed $B - V$ colors at the adopted temperature scale. By defining the color corrections in this way, we assume that the V -band bolometric corrections in the isochrones are good; $\Delta(V - I_C)$ is then taken as an adjustment to $BC(I_C)$ and applied to the calculated I magnitude of the isochrone (and equivalently for B and K_s).

The final step is to make the isochrone generate the right $B - V$, as illustrated in Figure 8, which plots the quantity $\Delta(B - V)$ against the isochrone effective temperature as derived from M_V . Here the smoothing was done in a moving box of width 200 K centered at intervals of 100 K, with adjacent pairs averaged.

Applying these corrections to the colors at each effective temperature defines the empirically calibrated isochrone for the Hyades, which is tabulated along with the color corrections in Table 2. The range of colors over which the correction was performed was limited by the following considerations. The Hyades turnoff is around $B - V = 0.2$ and the magnitude-limited Hipparcos

sample has few stars with $B - V > 1.3$. There are only a few stars with $B - V \leq 0.4$ so the correction is not well defined for the bluest colors. Therefore the most reliable color range for the corrections is $0.4 \leq B - V \leq 1.3$.

3. The color calibrations and metallicity

The process of empirically correcting the isochrones to match the Hyades necessarily erases many potential systematic errors in the input physics, metallicity scale, or the computation of isochrone colors. On the positive side, the isochrone now matches the shape of the Hyades main sequence to high precision and (on average) will yield the same effective temperature for all the various colors. We are thus prepared to derive more accurate distances and metallicities for other open clusters relative to the set of parameters assumed for the Hyades: $[\text{Fe}/\text{H}] = +0.13$, scaled solar abundances, $Y = 0.273$, $\alpha = 1.72$, $\langle(m - M)_0\rangle = 3.33$, etc.

There is, however, one more step before we can examine other clusters in detail, namely measuring the sensitivity of the isochrone luminosity to metallicity in the several colors. For this, we return to the individual color calibrations discussed above. The isochrone absolute magnitude M_V depends both on the luminosity of the isochrone in the theoretical plane and on the metallicity sensitivity of the adopted color-temperature relations. We have defined our empirical isochrone with respect to the Lejeune et al. (1998) color-temperature relation, and assume that the color corrections displayed in Table 2 are independent of metallicity. As a result, we will be measuring cluster distances on this system, and wish to compare this to the available alternatives.

In Figure 9, we illustrate the sensitivity of the main-sequence luminosity to metallicity for the three color-temperature relations we explored in § 2.2. The top panel plots the quantity

$$\Delta M_V(B - V) = \langle M_V([\text{Fe}/\text{H}]) - M_V([\text{Fe}/\text{H}] = +0.13) \rangle|_{B-V=0.4-1.0},$$

namely the average difference in absolute magnitude between the isochrone at any particular metallicity and that for $[\text{Fe}/\text{H}] = +0.13$, computed over the range $0.4 \leq (B - V) \leq 1.0$ at constant $B - V$. The points are at intervals of 0.1 in metallicity, from $[\text{Fe}/\text{H}] = -0.3$ to $+0.2$, and were computed by finding the average difference in M_V between the stellar values in Table 1 and the isochrone for that metallicity. The errors on the individual points are on the order of 0.01 mag. The values for the Hyades isochrone employing the Lejeune et al. (1998) color calibration are shown as filled points; the adjacent straight line is a linear least-squares fit to those points. The uncorrected Lejeune et al. (1998) and Alonso et al. (1995, 1996) color calibrations are displayed as open circles and triangles, respectively. The cross at $[\text{Fe}/\text{H}] = +0.13$ and $\Delta M_V = 0.0$ shows where the isochrones should go if they did not require empirical corrections to the color-temperature relation (i.e., if they matched the Hyades photometry *a priori*.)

The center and lower panels, respectively, plot the mean difference at fixed $V - I_C$ and $V - K_s$, computed over the same color interval in $B - V$.

The value of ΔM_V enters into the computation of cluster distances in several different ways. The slope of ΔM_V as a function of $[\text{Fe}/\text{H}]$ is a measure of distance errors that arise from errors in the assumed or measured metallicity of a cluster. The average slope over the range $-0.3 \leq [\text{Fe}/\text{H}] \leq +0.2$ is shown for the three representative color calibrations in Table 3 (the relationship is probably not precisely linear, having a somewhat steeper dependence on metallicity above $([\text{Fe}/\text{H}] = 0.0)$). All the color calibrations have nearly the same dependence on M_V at fixed $B - V$, with an average slope of 1.42 mag per dex in metallicity. The metallicity dependence is less steep at fixed $V - I_C$ or fixed $V - K_s$, and we note that the Alonso et al. (1995, 1996) calibration produces a markedly shallower metallicity dependence of the main-sequence luminosity in at constant $V - I_C$ than the other two calibrations.

While the various calibrations generally agree on the slope $d(\Delta M_V)/d[\text{Fe}/\text{H}]$, they have very different zero points. Since these calculations are all done for the same input physics and metallicity ($[\text{Fe}/\text{H}] = +0.13$), one would necessarily derive different distances to the Hyades using the various color calibrations. Forcing the distances to be the same for the various colors yields a photometric metallicity index for a cluster (e.g., Pinsonneault et al. 1998).

The last two columns of Table 3 display the value of $[\text{Fe}/\text{H}]$ that would be found for the various color calibrations, and the relative distance with respect to the assumed value of 3.33 for the Hyades.

4. Summary and discussion

In this paper, we have continued our effort to improve the calibration of main-sequence isochrones for use in determining the distances and metallicities of open clusters. We first identified a set of single Hyades stars with quality multicolor photometry, then verified that the isochrone luminosity-temperature relationship is in agreement with recent spectroscopic determinations. We then described a method for computing empirical corrections to the color-temperature relationship in the isochrone, and explored the sensitivity of the main-sequence luminosity to metallicity using several available color-temperature calibrations (see also VandenBerg and Clem 2003 for a parallel discussion of these issues). In subsequent papers, we will extend the empirical calibration to the lower main sequence via photometry in other clusters, determine the accuracy of photometric metallicity indicators, explore the effects of reddening, and derive ages for young systems via an analysis of the pre-main sequence / main sequence boundary.

The principal result of this paper was that the theoretical and spectroscopic (PSC) luminosity-temperature and luminosity-radius (i.e., L , $\log g$) relations are in fairly good agreement (Fig. 4) except for stars with $T_{\text{eff}} \gtrsim 5800$. The spectroscopic $\log g$ values, if real, would imply large errors in the model radii that would cause much larger differences in T_{eff} than seen (assuming that the mass/luminosity relationship is correct). They are also very unlikely given the similarity of the stars to the Sun, which should imply that the solar calibration produces reasonable radii for models of

Hyades stars. We have verified that the dispersion in all of the abundances measured by PSC is significantly reduced when the theoretical $\log g$ values are used in place of the spectroscopic ones (a more detailed analysis will appear in a future paper). Paulson et al. note similar trends with respect to temperature scales employed by Allende Prieto & Lambert (1999), which was based on a different set of theoretical isochrones. We suggest that there may be potential degeneracies between the spectroscopic temperature and gravity measurements. The spectroscopic T_{eff} and $\log g$ values are derived from Saha/Boltzmann considerations, and the fitting procedure used by PSC may compensate for errors in the ionization balance ($\log g$) by changing the excitation balance (T_{eff}). This idea could be tested by seeing what temperature scale resulted if the gravities were taken from the theoretical isochrones rather than treating them as a free parameter. In any case, the overall agreement in between a high-quality spectroscopic data set and the isochrones in the theoretical plane provides strong support for the hypothesis that the model temperature scale is reasonable.

The main goal of this effort is to reduce the theoretical and calibration uncertainties in the theoretical models using the constraints provided by the detailed morphology of the main sequence in various clusters. As outlined in the Introduction, however, we have long been interested in the discrepancy between the Hipparcos distances to clusters (ESA 1997) as subsequently analyzed by Robichon et al. (1999), van Leeuwen (1999), and others (see Pinsonneault et al. 1998; Soderblom et al. 1998). Various attempts to explain away the discrepancies as arising from, say, strange helium abundances (Belikov et al. 1998) or stellar activity (Percival, Salaris, & Kilkenney 2003), whether true or not (and worth exploring further), do raise an important issue: accurate calibration of isochrone physics and colors is an intrinsically difficult exercise, whereas parallaxes should be easier in principle (though perhaps not in practice).

In our view, the issue of the distance discrepancies arising from the 1997–1999 Hipparcos parallaxes is largely settled. Several papers have pointed out that small-scale errors in the parallaxes are effectively random zero-point errors over the whole sky, with an r.m.s. error of about 0.5 mas. The Hyades are less affected than other Hipparcos clusters because the Hyades subtend far larger than 1 degree, allowing the small-scale errors to average out Narayanan & Gould (1999b). The Pleiades are strongly affected because the bright stars are quite centrally concentrated (17 Hipparcos stars per square degree at the cluster center) and so they share the same zero-point error, which amounts to +1 mas at that point in the sky. Recently, Makarov (2002) has devised a method to recalculate parallaxes from the Hipparcos Intermediate Astrometric Data, free of the small angular scale correlated errors which are an artifact of the “great circle abscissa method” data reductions used in the ESA (1997) distances. His new Pleiades distance is in agreement with the distance from main-sequence fitting, and also with Pleiades distances derived from kinematic parallaxes (Narayanan & Gould 1999b) and new ground-based results (Gatewood, de Jonge, & Han 2000).

The work reported here was supported in part by the National Science Foundation, under grants AST-9731621 and AST-0206008 to the Ohio State University Research Foundation.

REFERENCES

- Allard, F., & Hauschildt P. H. 1995, *ApJ*, 445, 433
- Allende Prieto, C., & Lambert, D. W. 1999, *A&A*, 352, 555
- Alonso, A., Arribas, S., & Martínez-Roger, C. 1995, *A&A*, 297, 197
- Alonso, A., Arribas, S., & Martínez-Roger, C. 1996, *A&AS*, 117, 227
- Bahcall, J. N., Pinsonneault, M. H., & Basu, S. 2001, *ApJ*, 555, 990
- Basu, S., Pinsonneault, M. H., & Bahcall, J. N. 2000, *ApJ*, 529, 1084
- Belikov, A. N., Hirte, S., Meusinger, H., Piskunov, A. E., & Schilbach, E. 1998, *A&A*, 332, 575
- Bessel, M. S. 1979, *PASP*, 91, 589
- Bessel, M. S., & Weis, E. W. 1987, *PASP*, 99, 642
- Boesgaard, A. M., & Friel, E. D. 1990, *ApJ*, 351, 467
- de Bruijne, J. H. J., Hoogerwerf, R., & de Zeeuw, P. T. 2001, *A&A*, 367, 111
- Carney, B. W. 1982, *AJ*, 87, 1527
- Carney B. W., & Aaronson M. 1979, *AJ*, 84, 867
- Carpenter, J. M. 2001, *AJ*, 121, 2851
- Castellani, V., Degl’Innocenti, S., & Prada Moroni, P. G. 2001, *MNRAS*, 320, 66
- Crawford, D. L. 1975, *AJ*, 80, 955
- Eggen, O. J. 1948, *AJ*, 54, 35
- Eggen, O. J. 1975, *PASP*, 87, 107
- Eggen O. J., 1982, *ApJS*, 50, 221
- Elias, J. H., Frogel, J. A., Matthews, K., & Neugebauer, G. 1982, *AJ*, 87, 1029
- European Space Agency, 1997, *The Hipparcos and Tycho Catalogues* (ESA SP-1200) (Paris: ESA)
- Flower, P. J. 1996, *ApJ*, 469, 355
- Frogel, J. A., Persson, S. E., Aaronson, M., & Matthews, K. 1978, *ApJ*, 220, 75
- Gatewood, G., de Jonge, J. K., & Han, I. 2000, *ApJ*, 533, 938

- Grevesse, N., & Noels, A. 1993, in *Origin and Evolution of the Elements*, ed. M. Prantzos, E. Vangioni-Flam, & M. Cassé (Cambridge: Cambridge Univ. Press), 15
- Gratton, R. G., Carretta, E., & Castelli, F. 1996, *A&A*, 314, 191
- Grocholski, A. J., & Sarajedini, A. 2003, preprint (astro-ph/0307503)
- Johnson, H. L., & Knuckles, C. F. 1955, *ApJ*, 122, 209
- Johnson, H. L., MacArthur J. W., & Mitchell R. I. 1968, *ApJ*, 152, 465
- Johnson, H. L., Mitchell R. I., Iriarte B., & Wisniewski W. Z. 1966, *Comm. Lun. Plan. Lab.* 4, 99
- van Leeuwen, F. 1999, *A&A*, 341, 71
- van Leeuwen, F., & Evans, D. W. 1998, *A&AS*, 130, 157
- van Leeuwen, F., Alphenaar, P., & Meys, J. J. M. 1987, *A&A*, 67, 483
- Lejeune, T., Cuisinier, F., & Buser, R. 1998, *A&AS*, 130, 65
- Lejeune, T., & Schaerer, D. 2001, *A&A*, 366, 538
- Makarov, V. V. 2002, *AJ*, 124, 3299
- Mendoza, E. E. 1967, *Ton. Tac.* 4, 149
- Mermilliod, J.-C., Turon, C., Robichon, N., Arenou, F., & Lebreton, Y. 1997, in *Hipparcos Venice '97*, eds. B. Battick & M. A. C. Perryman, (Paris: ESA), 643
- Narayanan, V. K., & Gould, A. 1999a, *ApJ*, 515, 256
- Narayanan, V. K., & Gould, A. 1999b, *ApJ*, 523, 328
- Paulson, D. B., Sneden, C., & Cochran, W. D. 2003, *AJ*, 125, 3185 (PSC)
- Percival, S. M., Salaris, M., & Kilkenny, D. 2003, *A&A*, 400, 541
- Perryman, M. A. C., Brown, A. G. A., Lebreton, Y., Gómez, A., Turon, C., de Strobel, G. C., Mermilliod, J.-C., Robichon, N., Kovalevsky, J., & Crifo, F. 1998, *A&A*, 331, 81
- Persson, S. E., Murphy, D. C., Krzeminski, W., Roth, M., & Rieke, M. J. 1998, *AJ*, 116, 2475
- Pinsonneault, M. H., Stauffer, J., Soderblom, D. R., King, J. R., & Hanson, R. B. 1998, *ApJ*, 504, 170
- Pinsonneault, M. H., Terndrup, D. M., Hanson, R. B., & Stauffer, J. R. 2003, *ApJ*, in press, astro-ph/0307554 (Paper I)
- Robichon, N., Arenou, F., Mermilliod, J.-C., & Turon, C. 1999, *A&A*, 345, 471

- Robichon, N., Lebreton, Y., Turon, C., & Mermilliod, J.-C. 2000, in *Stellar Clusters and Associations: Convection, Rotation, and Dynamos*, eds. R. Pallavicini, G. Micela, and S. Sciortino, ASP Conf. Ser. 198, 141
- Saumon, D., Chabrier, G., & Van Horn, H. M. 1995, *ApJS*, 99, 713
- Soderblom, D. R., King, J. R., Hanson, R. B., Jones, B. F., Fischer, D., Stauffer, J. R., & Pinsonneault, M. H. 1998, *ApJ*, 504, 192
- Stauffer, J. R., Jones, B. F., Backman, D., Hartmann, L. W., Barrado y Nevascués, D., Pinsonneault, M. H., Terndrup, D. M., & Muench, A. 2003, *AJ*, 126, 833
- Stello, D., & Nissen, P. E. 2001, *A&A*, 374, 105
- Taylor, B. J. 1980, *AJ*, 85, 242
- Taylor, B. J., & Joner, M. D. 1985, *AJ*, 90, 479
- Terndrup, D. M., Stauffer, J. R., Pinsonneault, M. H., Sills, A., Yuan, Y., Jones, B. F., Fischer, D., & Krishnamurthi, A. 2000, *AJ*, 119, 1303
- Terndrup, D. M., Pinsonneault, M. H., Jeffries, R. D., Ford, A., Stauffer, J. R., & Sills, A. 2002, *ApJ*, 576, 950
- Ungren, A. R., & Weis, E. W. 1977, *ApJ*, 82, 978
- Ungren, A. R., Weis, E. W., & Hanson, R. B. 1985, *AJ*, 90, 2039
- Weis, E. W., Deluca, E. E., & Ungren, A. R. 1979, *PASP*, 91, 766
- Weis, E. W., & Hanson, R. B. 1988, *AJ*, 96, 148
- Weis, E. W. & Ungren, A. R. 1982, *PASP*, 94, 474
- Weiss, A., & Salaris, M. 1999, *A&A*, 346, 897
- VandenBerg, D. A., & Clem, J. L. 2003, *AJ*, 126, 778
- VandenBerg, D. A., Swenson, F. J., Rogers, F. J., Iglesias, C. A., & Alexander, D. R. 2000, *ApJ*, 532, 430
- Yi, S. K., Kim, Y.-C., & Demarque, P. 2003 *ApJS*, 144, 259

Table 1. Merged Photometry in the Hyades

Hipp	Other ID ^a	V	$\sigma(V)$	M_V	$\sigma(M_V)$	$B - V$	$\sigma(B - V)$	$V - I_C$	$\sigma(V - I_C)$	K_s	$\sigma(K_s)$	$V - K_s$	$\sigma(V - K_s)$	$J - K_s$	$\sigma(J - K_s)$	$H - K_s$	$\sigma(H - K_s)$
13806 vB 153		8.915	0.007	5.856	0.026	0.856	0.003	0.871	0.016	6.905	0.022	2.010	0.023	0.478	0.025	0.092	0.023
13834 vB 154		5.800	0.010	3.222	0.019	0.410	0.003	0.468	0.011	4.778	0.020	1.022	0.022	0.530	0.198	0.037	0.035
13976 G76-49		7.970	0.010	6.131	0.019	0.926	0.015	5.841	0.024	2.129	0.026	0.482	0.022	0.110	0.043
14976 LTT 11045		8.162	0.010	5.156	0.025	0.732	0.017	6.557	0.027	1.605	0.029	0.368	0.026	0.098	0.031
15563 G79-28		9.650	0.010	7.187	0.028	1.130	0.015	6.881	0.022	2.769	0.024	0.664	0.024	0.074	0.052
15720 LP 355-64		11.030	0.010	8.488	0.044	1.431	0.004	7.415	0.022	3.615	0.024	0.834	0.023	0.175	0.039
16529 vB 4		8.890	0.010	5.765	0.032	0.845	0.002	0.856	0.016	6.907	0.016	1.983	0.019	0.467	0.014	0.111	0.010
17766 G7-15		10.850	0.010	8.033	0.048	1.340	0.006	1.558	0.016	7.509	0.016	3.341	0.019	0.764	0.020	0.115	0.030
18170 vB 6		5.970	0.010	2.843	0.030	0.341	0.004	0.389	0.016	5.091	0.018	0.879	0.021	0.163	0.022	0.070	0.041
18322 L8, G7-34		10.120	0.010	6.779	0.053	1.070	0.008	1.124	0.020	7.574	0.021	2.546	0.023	0.607	0.020	0.118	0.024
18327 vB 7		8.987	0.010	5.927	0.037	0.895	0.002	0.899	0.011	6.910	0.021	2.077	0.023	0.503	0.019	0.128	0.027
19082 L12		11.415	0.007	8.007	0.071	1.348	0.004	1.577	0.020	8.107	0.026	3.308	0.027	0.782	0.034	0.152	0.031
19098 L10		9.310	0.015	6.032	0.044	0.893	0.005	0.896	0.016	7.254	0.022	2.056	0.027	0.494	0.030	0.111	0.039
19148 vB 10		7.850	0.006	4.449	0.036	0.591	0.003	0.607	0.016	6.446	0.024	1.404	0.025	0.307	0.025	0.086	0.038
19207 L15		10.485	0.007	7.163	0.051	1.180	0.008	1.268	0.020	7.665	0.021	2.820	0.022	0.700	0.028	0.146	0.051
19263 BD+16:558		9.940	0.010	6.628	0.044	1.005	0.012	7.511	0.026	2.429	0.028	0.594	0.027	0.088	0.044
19316 L14		11.277	0.006	7.871	0.074	1.328	0.003	1.558	0.016	8.046	0.027	3.231	0.028	0.772	0.033	0.125	0.045
19441 BD+8:642		10.100	0.010	7.361	0.045	1.198	0.006	1.278	0.016	7.264	0.020	2.836	0.022	0.648	0.016	0.076	0.021
19781 vB 17		8.460	0.007	4.947	0.046	0.696	0.004	0.720	0.011	6.805	0.009	1.655	0.012	0.391	0.025	0.091	0.021
19786 vB 18		8.060	0.007	4.735	0.046	0.640	0.002	0.677	0.011	6.535	0.009	1.525	0.012	0.322	0.013	0.073	0.017
19789 vB 16		7.057	0.006	3.283	0.037	0.420	0.003	0.451	0.016	6.005	0.021	1.052	0.022	0.202	0.017	0.077	0.023
19793 vB 15		8.067	0.020	4.820	0.039	0.658	0.003	0.692	0.011	6.516	0.020	1.551	0.028	0.361	0.023	0.083	0.018
19796 vB 19		7.130	0.010	3.846	0.038	0.512	0.003	0.560	0.016	5.862	0.017	1.268	0.020	0.266	0.013	0.053	0.017
19808 vA 68		10.725	0.050	7.424	0.085	1.205	0.005	1.288	0.011	7.680	0.009	3.045	0.050	0.757	0.025	0.140	0.013
19834 vA 72		11.557	0.015	8.197	0.115	1.373	0.006	1.630	0.011	8.141	0.009	3.416	0.018	0.819	0.024	0.166	0.021
19877 vB 20		6.321	0.007	2.987	0.033	0.399	0.004	0.459	0.016	5.340	0.009	0.981	0.011	0.188	0.030	0.075	0.020
19934 vB 21		9.147	0.006	5.625	0.043	0.814	0.002	0.825	0.011	7.275	0.022	1.872	0.023	0.426	0.020	0.101	0.020
20130 vB 26		8.625	0.007	5.323	0.045	0.743	0.003	0.762	0.016	6.921	0.016	1.704	0.017	0.365	0.012	0.079	0.012
20146 vB 27		8.464	0.015	5.138	0.045	0.717	0.004	0.749	0.011	6.793	0.009	1.671	0.018	0.347	0.017	0.085	0.015
20219 vB 30		5.591	0.007	2.331	0.035	0.279	0.003	0.338	0.011	4.861	0.009	0.730	0.011	0.149	0.033	0.074	0.045
20237 vB 31		7.467	0.006	4.200	0.034	0.566	0.004	0.622	0.016	6.124	0.018	1.343	0.019	0.271	0.013	0.060	0.024
20261 vB 33		5.266	0.007	1.879	0.037	0.223	0.003	0.272	0.011	4.689	0.016	0.577	0.017	0.061	0.046	0.284	0.282
20349 vB 35		6.801	0.006	3.329	0.039	0.436	0.004	0.513	0.016	5.762	0.020	1.039	0.021	0.208	0.017	0.024	0.015
20350 vB 36		6.807	0.006	3.457	0.035	0.441	0.004	0.498	0.016	5.746	0.016	1.061	0.017	0.198	0.015	0.048	0.010
20357 vB 37		6.611	0.006	3.160	0.038	0.406	0.004	0.475	0.016	5.610	0.009	1.000	0.011	0.189	0.019	0.071	0.031
20480 vB 42		8.854	0.010	5.338	0.048	0.758	0.002	0.786	0.016	7.121	0.016	1.733	0.019	0.410	0.011	0.076	0.011
20485 vB 173		10.499	0.020	7.485	0.072	1.237	0.003	1.368	0.011	7.481	0.009	3.018	0.022	0.669	0.019	0.153	0.039

Table 1—Continued

Hipp	Other ID ^a	V	$\sigma(V)$	M_V	$\sigma(M_V)$	$B - V$	$\sigma(B - V)$	$V - I_C$	$\sigma(V - I_C)$	K_s	$\sigma(K_s)$	$V - K_s$	$\sigma(V - K_s)$	$J - K_s$	$\sigma(J - K_s)$	$H - K_s$	$\sigma(H - K_s)$
20491	vB 44	7.185	0.007	3.541	0.042	0.452	0.003	0.521	0.016	6.092	0.016	1.093	0.017	0.193	0.015	0.049	0.010
20492	vB 46	9.117	0.006	5.731	0.054	0.865	0.003	0.871	0.011	7.153	0.009	1.964	0.011	0.466	0.011	0.110	0.016
20527	vA 294	10.901	0.020	7.673	0.075	1.288	0.002	1.465	0.009	7.755	0.009	3.146	0.022	0.720	0.021	0.139	0.021
20542	vB 47	4.800	0.010	1.526	0.038	0.156	0.003	0.179	0.016	4.400	0.010	0.400	0.014	0.147	0.268	0.099	0.210
20557	vB 48	7.141	0.006	4.010	0.037	0.521	0.004	0.576	0.016	5.871	0.020	1.270	0.021	0.277	0.020	0.075	0.025
20563	vB 174	9.995	0.005	6.740	0.064	1.058	0.003	1.101	0.011	7.525	0.009	2.470	0.011	0.630	0.013	0.112	0.020
20567	vB 51	6.971	0.006	3.461	0.047	0.443	0.004	0.517	0.016	5.880	0.009	1.090	0.011	0.167	0.014	0.043	0.016
20635	vB 54	4.220	0.010	0.846	0.034	0.137	0.003	0.163	0.011	4.077	0.470	0.404	0.470	0.016	0.549	-0.013	0.522
20641	vB 55	5.281	0.007	2.048	0.033	0.248	0.003	0.311	0.016	4.607	0.016	0.674	0.017	0.479	0.250	0.314	0.230
20741	vB 64	8.120	0.007	4.850	0.046	0.659	0.003	0.694	0.011	6.554	0.010	1.566	0.012	0.346	0.021	0.096	0.036
20762	vA 407	10.475	0.007	7.118	0.062	1.146	0.002	1.230	0.016	7.743	0.010	2.732	0.013	0.637	0.046	0.122	0.037
20815	vB 65	7.421	0.006	4.054	0.039	0.536	0.004	0.584	0.011	6.129	0.010	1.292	0.012	0.234	0.020	0.033	0.028
20826	vB 66	7.507	0.010	4.252	0.042	0.556	0.003	0.607	0.016	6.158	0.017	1.349	0.020	0.266	0.011	0.064	0.027
20827	vB 179	9.502	0.015	6.063	0.060	0.931	0.003	0.959	0.016	7.346	0.010	2.156	0.018	0.546	0.028	0.108	0.021
20842	vB 67	5.720	0.010	2.251	0.039	0.271	0.004	0.303	0.016	5.055	0.018	0.665	0.021	0.108	0.011	0.048	0.014
20894	vB 72	3.410	0.010	0.146	0.036	0.179	0.003	0.210	0.011	2.896	0.011	0.514	0.015	0.114	0.335	0.117	0.339
20899	vB 73	7.847	0.010	4.521	0.041	0.609	0.003	0.626	0.011	6.398	0.010	1.449	0.014	0.292	0.021	0.058	0.025
20948	vB 78	6.914	0.010	3.609	0.042	0.451	0.002	0.516	0.009	5.791	0.010	1.123	0.014	0.236	0.029	0.078	0.034
20949	vB 76	9.206	0.030	5.394	0.057	0.764	0.002	0.768	0.011	7.462	0.029	1.744	0.042	0.382	0.028	0.072	0.034
20951	vB 79	8.955	0.007	5.697	0.055	0.831	0.002	0.827	0.009	7.056	0.010	1.899	0.012	0.469	0.029	0.094	0.029
20978	vB 180	9.090	0.030	5.807	0.051	0.854	0.003	0.879	0.016	7.120	0.010	1.970	0.032	0.480	0.025	0.072	0.030
21029	vB 82	4.786	0.007	1.484	0.034	0.171	0.002	0.187	0.016	4.366	0.011	0.420	0.013	0.409	0.226	0.313	0.203
21036	vB 84	5.406	0.007	2.159	0.036	0.262	0.003	0.311	0.016	4.732	0.009	0.674	0.012	0.473	0.312	0.086	0.034
21066	vB 86	7.045	0.007	3.746	0.040	0.465	0.004	0.529	0.016	5.893	0.020	1.152	0.021	0.246	0.032	0.051	0.028
21099	vB 87	8.590	0.010	5.276	0.048	0.742	0.004	0.747	0.016	6.884	0.022	1.706	0.024	0.375	0.020	0.081	0.063
21112	vB 88	7.775	0.007	4.238	0.049	0.539	0.004	0.591	0.016	6.464	0.010	1.311	0.012	0.278	0.019	0.117	0.051
21138	vB 191	11.055	0.015	7.678	0.121	1.293	0.003	1.471	0.016	7.895	0.024	3.160	0.028	0.772	0.024	0.139	0.020
21152	vB 90	6.391	0.010	3.280	0.038	0.413	0.004	0.490	0.016	5.333	0.021	1.058	0.023	0.260	0.023	0.052	0.020
21256	L66	10.687	0.015	7.515	0.060	1.236	0.004	1.362	0.016	7.686	0.026	3.001	0.030	0.741	0.026	0.149	0.025
21261	L65	10.737	0.015	7.410	0.070	1.198	0.003	1.317	0.016	7.820	0.022	2.917	0.027	0.715	0.021	0.157	0.021
21267	vB 94	6.620	0.010	3.319	0.044	0.431	0.004	0.482	0.016	5.558	0.022	1.062	0.024	0.201	0.020	0.054	0.020
21317	vB 97	7.920	0.010	4.633	0.046	0.635	0.004	0.638	0.016	6.454	0.010	1.466	0.014	0.300	0.022	0.107	0.034
21637	vB 105	7.527	0.010	4.364	0.036	0.576	0.003	0.607	0.016	6.162	0.022	1.365	0.024	0.292	0.021	0.036	0.018
21723	L80	10.043	0.015	6.847	0.066	1.072	0.004	1.136	0.016	7.480	0.022	2.563	0.027	0.609	0.021	0.122	0.024
21741	vB 109	9.397	0.007	5.495	0.058	0.812	0.002	0.816	0.011	7.565	0.020	1.832	0.021	0.431	0.017	0.066	0.015
22044	vB 111	5.401	0.007	2.184	0.045	0.251	0.002	0.296	0.016	4.733	0.020	0.668	0.021	0.229	0.248	-0.001	0.026
22177	L119	10.910	0.010	7.623	0.091	1.276	0.004	1.404	0.020	7.826	0.020	3.084	0.022	0.744	0.015	0.158	0.035

Table 1—Continued

Hipp	Other ID ^a	V	$\sigma(V)$	M_V	$\sigma(M_V)$	$B - V$	$\sigma(B - V)$	$V - I_C$	$\sigma(V - I_C)$	K_s	$\sigma(K_s)$	$V - K_s$	$\sigma(V - K_s)$	$J - K_s$	$\sigma(J - K_s)$	$H - K_s$	$\sigma(H - K_s)$
22253	L93	10.687	0.006	6.999	0.066	1.112	0.003	1.184	0.016	7.995	0.033	2.692	0.034	0.695	0.045	0.139	0.036
22380	vB 116	8.990	0.010	5.592	0.062	0.828	0.002	0.846	0.011	7.059	0.026	1.931	0.028	0.427	0.027	0.082	0.028
22422	vB 118	7.737	0.010	4.327	0.048	0.580	0.003	0.607	0.016	6.355	0.016	1.382	0.019	0.287	0.015	0.078	0.017
22566	vB 143	7.895	0.007	3.975	0.059	0.526	0.003	0.568	0.016	6.671	0.021	1.224	0.022	0.259	0.031	0.015	0.017
22654	L98	10.290	0.010	6.704	0.081	1.070	0.008	1.114	0.020	7.742	0.020	2.548	0.022	0.633	0.025	0.143	0.026
22850	vB 126	6.371	0.007	2.378	0.056	0.291	0.004	0.350	0.016	5.623	0.018	0.748	0.019	0.113	0.022	0.035	0.014
23069	vB 127	8.890	0.007	5.175	0.073	0.737	0.002	0.724	0.016	7.235	0.017	1.655	0.018	0.334	0.013	0.033	0.054
23214	vB 128	6.755	0.007	3.593	0.039	0.450	0.004	0.498	0.016	5.645	0.016	1.110	0.017	0.211	0.018	0.049	0.030
23312	BD+04:810	9.722	0.010	6.159	0.074	0.957	0.047	7.589	0.026	2.133	0.028	0.524	0.026	0.103	0.039
23497	vB 129	4.640	0.010	1.041	0.045	0.150	0.003	0.195	0.016	4.245	0.021	0.395	0.023	0.088	0.320	0.130	0.209
23750	BD+17:841	8.820	0.010	5.197	0.061	0.730	0.015	7.128	0.017	1.692	0.020	0.386	0.011	0.088	0.044
24116	BD+20:897	7.852	0.010	3.314	0.084	0.445	0.015	6.821	0.017	1.031	0.020	0.214	0.021	0.059	0.013
24923	BD+11:772	9.042	0.010	5.292	0.088	0.765	0.029	7.302	0.021	1.740	0.023	0.394	0.021	0.095	0.061
26382	vB 168	5.541	0.006	2.014	0.072	0.222	0.004	0.257	0.016	4.935	0.025	0.606	0.026	0.127	0.027	0.067	0.031
28356	BD+02:1102	7.780	0.010	3.422	0.204	0.461	0.015	6.685	0.020	1.095	0.022	0.260	0.020	0.040	0.026

^aG = Giclas. L = Leiden (Pels). LP = Luyten Palomar. LTT = Luyten Two Tenths. vB = van Buren. vA = van Altena.

Table 2. Empirically Calibrated Hyades Isochrone

$T_{\text{eff}}(\text{K})$	M/M_{\odot}	M_V	$B - V$	$V - I_C$	$V - K_s$	$\Delta(B - V)$	$\Delta(V - I_C)$	$\Delta(V - K_s)$
8280	2.235	1.10	0.139	0.162	0.374	0.027	0.053	0.014
8276	2.214	1.15	0.141	0.164	0.377	0.027	0.054	0.015
8270	2.192	1.20	0.143	0.167	0.380	0.027	0.054	0.016
8262	2.170	1.25	0.146	0.170	0.384	0.027	0.055	0.017
8251	2.149	1.30	0.149	0.174	0.390	0.027	0.056	0.018
8236	2.127	1.35	0.153	0.178	0.396	0.027	0.057	0.019
8215	2.105	1.40	0.158	0.184	0.409	0.027	0.057	0.024
8189	2.083	1.45	0.163	0.191	0.425	0.027	0.058	0.032
8158	2.062	1.50	0.170	0.199	0.443	0.026	0.058	0.040
8124	2.040	1.55	0.176	0.208	0.461	0.026	0.059	0.046
8088	2.019	1.60	0.183	0.218	0.479	0.025	0.059	0.051
8052	1.997	1.65	0.190	0.228	0.496	0.024	0.059	0.056
8015	1.976	1.70	0.197	0.238	0.514	0.023	0.059	0.061
7978	1.955	1.75	0.204	0.247	0.532	0.021	0.058	0.065
7942	1.934	1.80	0.210	0.256	0.549	0.020	0.057	0.069
7905	1.914	1.85	0.216	0.265	0.566	0.018	0.055	0.072
7868	1.893	1.90	0.223	0.273	0.582	0.016	0.053	0.075
7830	1.873	1.95	0.229	0.281	0.599	0.014	0.051	0.077
7790	1.854	2.00	0.235	0.287	0.614	0.012	0.049	0.079
7749	1.834	2.05	0.242	0.294	0.630	0.010	0.048	0.081
7706	1.815	2.10	0.249	0.300	0.646	0.008	0.046	0.082
7662	1.796	2.15	0.256	0.305	0.662	0.006	0.045	0.083
7616	1.776	2.20	0.263	0.310	0.678	0.004	0.045	0.084
7568	1.757	2.25	0.271	0.315	0.694	0.002	0.044	0.085
7520	1.738	2.30	0.279	0.320	0.709	0.001	0.044	0.085
7471	1.720	2.35	0.286	0.324	0.724	−0.001	0.043	0.085
7423	1.702	2.40	0.294	0.329	0.739	−0.002	0.043	0.085
7375	1.685	2.45	0.301	0.334	0.754	−0.002	0.042	0.085
7327	1.667	2.50	0.308	0.341	0.769	−0.003	0.040	0.084
7280	1.651	2.55	0.315	0.348	0.784	−0.003	0.039	0.084
7232	1.634	2.60	0.322	0.356	0.799	−0.003	0.038	0.083
7186	1.618	2.65	0.329	0.365	0.814	−0.004	0.037	0.083
7139	1.603	2.70	0.336	0.375	0.829	−0.004	0.036	0.083
7092	1.587	2.75	0.343	0.386	0.845	−0.004	0.035	0.082
7046	1.571	2.80	0.350	0.397	0.860	−0.004	0.035	0.081
7001	1.556	2.85	0.358	0.408	0.876	−0.004	0.034	0.080
6956	1.541	2.90	0.365	0.419	0.892	−0.005	0.033	0.079
6913	1.526	2.95	0.372	0.429	0.909	−0.005	0.032	0.077
6871	1.511	3.00	0.379	0.439	0.926	−0.006	0.031	0.075
6830	1.496	3.05	0.387	0.449	0.943	−0.006	0.031	0.072
6792	1.482	3.10	0.394	0.457	0.961	−0.007	0.029	0.069
6754	1.468	3.15	0.401	0.465	0.980	−0.008	0.028	0.065

Table 2—Continued

$T_{\text{eff}}(\text{K})$	M/M_{\odot}	M_V	$B - V$	$V - I_C$	$V - K_s$	$\Delta(B - V)$	$\Delta(V - I_C)$	$\Delta(V - K_s)$
6718	1.454	3.20	0.408	0.472	0.998	−0.009	0.027	0.062
6684	1.440	3.25	0.415	0.479	1.016	−0.010	0.025	0.057
6650	1.426	3.30	0.422	0.485	1.034	−0.010	0.023	0.052
6618	1.413	3.35	0.429	0.491	1.051	−0.011	0.021	0.047
6586	1.400	3.40	0.436	0.497	1.068	−0.012	0.020	0.041
6556	1.387	3.45	0.443	0.503	1.084	−0.012	0.018	0.034
6525	1.375	3.50	0.450	0.508	1.099	−0.012	0.016	0.028
6496	1.362	3.55	0.457	0.514	1.116	−0.012	0.015	0.023
6466	1.350	3.60	0.464	0.521	1.132	−0.012	0.014	0.017
6438	1.338	3.65	0.471	0.527	1.148	−0.012	0.013	0.012
6409	1.326	3.70	0.478	0.534	1.164	−0.011	0.012	0.008
6382	1.315	3.75	0.485	0.540	1.181	−0.010	0.011	0.003
6354	1.303	3.80	0.493	0.546	1.197	−0.010	0.010	0.000
6327	1.292	3.85	0.500	0.552	1.214	−0.009	0.010	−0.003
6300	1.281	3.90	0.507	0.558	1.231	−0.007	0.008	−0.007
6273	1.269	3.95	0.515	0.564	1.247	−0.006	0.007	−0.010
6246	1.259	4.00	0.522	0.570	1.264	−0.005	0.006	−0.013
6220	1.248	4.05	0.530	0.576	1.281	−0.003	0.005	−0.015
6194	1.237	4.10	0.538	0.582	1.299	−0.002	0.004	−0.017
6167	1.227	4.15	0.545	0.589	1.318	0.000	0.003	−0.018
6141	1.217	4.20	0.553	0.595	1.337	0.002	0.002	−0.019
6115	1.207	4.25	0.561	0.601	1.357	0.003	0.001	−0.018
6089	1.197	4.30	0.569	0.608	1.377	0.005	0.000	−0.018
6063	1.187	4.35	0.577	0.614	1.396	0.007	0.000	−0.018
6036	1.177	4.40	0.585	0.621	1.414	0.009	−0.001	−0.018
6010	1.167	4.45	0.594	0.628	1.430	0.010	−0.001	−0.019
5982	1.157	4.50	0.602	0.636	1.445	0.012	−0.002	−0.020
5955	1.147	4.55	0.611	0.643	1.460	0.013	−0.002	−0.021
5927	1.137	4.60	0.619	0.651	1.474	0.015	−0.003	−0.021
5899	1.128	4.65	0.628	0.658	1.488	0.016	−0.003	−0.021
5871	1.118	4.70	0.637	0.666	1.503	0.018	−0.004	−0.020
5843	1.108	4.75	0.646	0.674	1.519	0.019	−0.005	−0.019
5815	1.099	4.80	0.654	0.682	1.535	0.020	−0.005	−0.018
5787	1.090	4.85	0.663	0.690	1.551	0.021	−0.006	−0.018
5760	1.080	4.90	0.672	0.698	1.568	0.021	−0.006	−0.018
5732	1.071	4.95	0.681	0.706	1.586	0.022	−0.007	−0.018
5704	1.062	5.00	0.690	0.715	1.605	0.022	−0.007	−0.019
5676	1.054	5.05	0.700	0.724	1.625	0.023	−0.007	−0.020
5648	1.045	5.10	0.709	0.733	1.646	0.023	−0.007	−0.021
5620	1.036	5.15	0.719	0.742	1.667	0.023	−0.007	−0.024
5591	1.028	5.20	0.729	0.751	1.689	0.023	−0.008	−0.026
5561	1.020	5.25	0.739	0.760	1.711	0.023	−0.008	−0.029

Table 2—Continued

$T_{\text{eff}}(\text{K})$	M/M_{\odot}	M_V	$B - V$	$V - I_C$	$V - K_s$	$\Delta(B - V)$	$\Delta(V - I_C)$	$\Delta(V - K_s)$
5532	1.012	5.30	0.749	0.768	1.733	0.023	−0.008	−0.032
5502	1.003	5.35	0.759	0.777	1.756	0.023	−0.008	−0.036
5471	0.995	5.40	0.769	0.786	1.779	0.023	−0.008	−0.040
5440	0.988	5.45	0.780	0.795	1.802	0.023	−0.008	−0.044
5408	0.980	5.50	0.790	0.804	1.824	0.022	−0.007	−0.048
5377	0.972	5.55	0.801	0.813	1.847	0.021	−0.006	−0.052
5344	0.964	5.60	0.811	0.823	1.870	0.020	−0.004	−0.056
5312	0.957	5.65	0.822	0.833	1.893	0.019	−0.002	−0.059
5279	0.949	5.70	0.833	0.843	1.917	0.017	0.001	−0.061
5245	0.942	5.75	0.844	0.854	1.941	0.015	0.004	−0.061
5212	0.934	5.80	0.854	0.866	1.966	0.013	0.008	−0.060
5179	0.927	5.85	0.865	0.877	1.993	0.011	0.011	−0.058
5146	0.920	5.90	0.876	0.888	2.019	0.008	0.015	−0.056
5113	0.913	5.95	0.887	0.900	2.046	0.005	0.018	−0.054
5080	0.906	6.00	0.898	0.911	2.072	0.002	0.020	−0.053
5048	0.899	6.05	0.908	0.921	2.098	−0.001	0.023	−0.052
5017	0.892	6.10	0.919	0.933	2.125	−0.004	0.025	−0.053
4986	0.886	6.15	0.930	0.944	2.151	−0.006	0.027	−0.053
4956	0.879	6.20	0.941	0.955	2.179	−0.009	0.029	−0.054
4927	0.873	6.25	0.952	0.966	2.206	−0.012	0.031	−0.056
4898	0.867	6.30	0.963	0.978	2.235	−0.014	0.032	−0.058
4870	0.861	6.35	0.974	0.990	2.263	−0.016	0.033	−0.060
4843	0.855	6.40	0.985	1.003	2.293	−0.019	0.034	−0.061
4816	0.850	6.45	0.996	1.016	2.322	−0.021	0.034	−0.063
4790	0.844	6.50	1.007	1.030	2.353	−0.022	0.035	−0.064
4765	0.839	6.55	1.019	1.044	2.384	−0.024	0.035	−0.065
4740	0.833	6.60	1.030	1.059	2.415	−0.026	0.034	−0.065
4715	0.828	6.65	1.041	1.074	2.446	−0.027	0.034	−0.065
4691	0.823	6.70	1.053	1.090	2.478	−0.028	0.033	−0.065
4667	0.818	6.75	1.064	1.106	2.510	−0.029	0.032	−0.064
4644	0.813	6.80	1.076	1.122	2.542	−0.030	0.031	−0.062
4620	0.808	6.85	1.087	1.138	2.574	−0.031	0.030	−0.060
4597	0.803	6.90	1.099	1.155	2.606	−0.031	0.028	−0.057
4573	0.798	6.95	1.110	1.171	2.638	−0.031	0.027	−0.054
4550	0.793	7.00	1.121	1.187	2.670	−0.031	0.025	−0.050
4527	0.788	7.05	1.132	1.203	2.702	−0.031	0.024	−0.045
4503	0.783	7.10	1.144	1.219	2.734	−0.030	0.023	−0.039
4480	0.778	7.15	1.155	1.235	2.765	−0.029	0.021	−0.033
4457	0.773	7.20	1.167	1.252	2.798	−0.027	0.021	−0.026
4435	0.768	7.25	1.178	1.268	2.830	−0.025	0.021	−0.018
4412	0.763	7.30	1.190	1.285	2.862	−0.022	0.021	−0.009
4390	0.757	7.35	1.202	1.302	2.894	−0.019	0.022	0.000

Table 2—Continued

$T_{\text{eff}}(\text{K})$	M/M_{\odot}	M_V	$B - V$	$V - I_C$	$V - K_s$	$\Delta(B - V)$	$\Delta(V - I_C)$	$\Delta(V - K_s)$
4368	0.752	7.40	1.214	1.320	2.927	−0.015	0.023	0.011
4347	0.747	7.45	1.226	1.339	2.961	−0.010	0.026	0.023
4326	0.742	7.50	1.238	1.357	2.994	−0.006	0.029	0.035
4305	0.737	7.55	1.250	1.378	3.028	−0.001	0.034	0.048
4285	0.731	7.60	1.262	1.400	3.061	0.004	0.039	0.061
4265	0.726	7.65	1.274	1.422	3.093	0.009	0.045	0.073
4246	0.721	7.70	1.285	1.445	3.123	0.014	0.052	0.083
4228	0.715	7.75	1.296	1.468	3.151	0.018	0.059	0.092
4209	0.710	7.80	1.307	1.493	3.177	0.023	0.066	0.098
4192	0.705	7.85	1.317	1.517	3.199	0.028	0.074	0.102
4175	0.700	7.90	1.328	1.541	3.221	0.032	0.080	0.104
4158	0.694	7.95	1.338	1.562	3.244	0.037	0.084	0.108
4142	0.689	8.00	1.348	1.580	3.270	0.042	0.085	0.115
4126	0.684	8.05	1.358	1.596	3.303	0.046	0.085	0.129
4110	0.678	8.10	1.368	1.616	3.341	0.051	0.087	0.148
4095	0.673	8.15	1.377	1.643	3.385	0.055	0.098	0.172
4080	0.668	8.20	1.386	1.676	3.431	0.059	0.115	0.200
4066	0.663	8.25	1.395	1.711	3.477	0.063	0.135	0.226
4051	0.657	8.30	1.403	1.756	3.508	0.067	0.165	0.238
4038	0.652	8.35	1.411	1.801	3.532	0.071	0.195	0.242
4024	0.647	8.40	1.419	1.844	3.554	0.074	0.224	0.244
4010	0.642	8.45	1.426	1.882	3.586	0.077	0.250	0.256
3997	0.636	8.50	1.432	1.927	3.624	0.080	0.282	0.274

Table 3. Metallicity sensitivity of isochrone

Color calibration	$B - V$	$V - I_C$	$V - K_s$	[M/H]	$\Delta(m - M)_V$
Lejeune et al. (1998)	1.402	0.927	0.804	+0.142	0.00
Uncorrected Lejeune	1.432	0.903	0.767	−0.019	−0.10
Alonso et al. (1995,1996)	1.421	0.727	0.758	+0.036	−0.40

Fig. 1.— Comparison of Hyades single-star photometry (points) in M_V , $B - V$ to a theoretical isochrone (solid line) for $[\text{Fe}/\text{H}] = +0.13$, age 550 Myr, which uses the Lejeune et al. (1998) color calibration. Shown as a long-dashed line is the same isochrone, but employing the uncalibrated Lejeune et al. (1998) color-temperature relation; the short-dashed line is the equivalent for the Alonso et al. (1995, 1996) calibration. Error bars are shown for all points, even though these are typically smaller than the points themselves.

Fig. 2.— Same as Figure 1, but in M_V , $V - I_C$.

Fig. 3.— Same as Figure 1, but in M_V , $V - K_s$.

Fig. 4.— Differences in effective temperature (top panel) and gravity (lower panel), in the sense of (isochrone – spectra), between spectroscopically determined parameters (Paulson, Sneden, & Cochran 2003) and the Hyades isochrone at fixed M_V . While the agreement is good throughout most of the temperature range, the spectroscopic solution yields hotter temperatures and lower gravities than the isochrone. Filled circles are for the stars used to calibrate the empirical color corrections, while the triangles show other stars in the Paulson et al. sample not included in our Table 1.

Fig. 5.— Upper panel: iron abundances for Hyades stars from Paulson, Sneden, & Cochran (2003) as a function of isochrone effective temperature. Lower panel: rescaled iron abundances using the isochrone temperatures and gravities, as described in the text. Symbols are the same as in Figure 4.

Fig. 6.— Differences between T_{eff} estimates from $B - V$ and $V - I_C$ (top panel), from $B - V$ and $V - K_s$ (center panel), and from $V - I_C$ and $V - K_s$ (lower panel) for the Lejeune color calibration as a function of T_{eff} estimated from the isochrone M_V .

Fig. 7.— Method of constructing color corrections to the theoretical isochrone. The points in the top panel show the difference in $V - I_C$ between the isochrone (Figure 1) and the data at the isochrone color. The sense of the difference is (star – isochrone). The smoothed line was constructed by computing a weighted average difference in a moving window containing 3–5 stars. The lower panels is the equivalent for $V - K_s$. As discussed in the text, the corrections are applied to the isochrone colors holding M_V fixed.

Fig. 8.— Color correction in $B - V$.

Fig. 9.— Sensitivity of the isochrone luminosity to metallicity, for the several color calibrations discussed in the text.

















